Redondo Submarine Canyon: Formation, Geology, and Sediment Dynamics

A Comprehensive Report on the Underwater Canyon off Redondo Beach Harbor

---

Executive Summary

Redondo Canyon is a spectacular submarine canyon in the Santa Monica Bay that begins just offshore from Redondo Beach Harbor, California, and extends approximately 15.6 kilometers to depths exceeding 800 meters. This active geological feature was carved primarily by the ancestral San Gabriel River during Pleistocene glacial periods when sea levels were up to 122 meters lower than today. The canyon remains dynamically active, transporting sediment through turbidity currents and other mass-wasting processes, and is clearly visible on modern bathymetric imagery including Google Earth. This report examines the canyon's formation mechanisms, morphology, sediment transport systems, and continuing geological activity.

Table of Contents

1. Introduction and Location
2. Canyon Morphology and Dimensions
3. Geological Formation History
4. Pleistocene Glacial Periods and Sea Level Change
5. The San Gabriel River Connection
6. Submarine Canyon Formation Mechanisms
7. Turbidity Currents and Sediment Transport
8. Littoral Drift and Modern Sediment Supply
9. The Redondo Submarine Fan
10. Modern Activity and Holocene Dynamics
11. California Continental Borderland Context
12. Observing Redondo Canyon Today
13. Comparison with Other Southern California Canyons
14. Scientific Research and Discoveries
15. Ecological Significance
16. Conclusions

Introduction and Location

Redondo Canyon is a major submarine canyon system located off the coast of Redondo Beach in the South Bay region of Los Angeles County, California. The canyon is part of the Santa Monica Bay submarine canyon system within the larger California Continental Borderland—a geologically complex region of basins, ridges, and fault-controlled topography extending from Point Conception to the United States-Mexico border.

Geographic Position

- **Canyon Head Location**: Approximately 100-200 meters offshore from Redondo Beach Harbor
- **Latitude/Longitude**: The canyon head begins at approximately 33°50'N, 118°24'W
- **Trend**: West-southwest to south-southwest
- **Termination**: San Pedro Basin at depths exceeding 800 meters
- **Total Length**: Approximately 15.6 kilometers from head to fan terminus

The canyon is uniquely positioned among Southern California submarine canyons because its head comes remarkably close to shore, beginning at depths as shallow as 10 meters (33 feet), making it accessible to recreational divers and visible on modern seafloor mapping systems.



Canyon Morphology and Dimensions

Canyon Head Characteristics

The canyon head—the uppermost, shallowest portion—displays dramatic relief despite its proximity to shore:

- **Initial Depth**: 10-30 meters (33-100 feet) below sea level
- **Steep Gradient**: The canyon wall drops at approximately 30° angle beginning around 10-12 meters depth
- **Rapid Descent**: Within a short horizontal distance, the canyon floor plunges from 30 meters to 60-80 meters, then exceeds 100 meters depth before descending "to the abyss"
- **Width at Head**: Relatively narrow, approximately 100-200 meters wide
- **Location**: Directly offshore from King Harbor/Redondo Beach Marina

This unusually shallow canyon head is critical to understanding the canyon's formation and continued activity. The proximity to shore allows the canyon to intercept littoral (nearshore) sediment transported by longshore currents, wave action, and rip currents.

Mid-Canyon Profile

As the canyon extends seaward and deepens:

- **Axis Orientation**: Remarkably straight, trending west-southwest
- **Width**: Expands to 800+ meters in some sections
- **Incision Depth**: Approximately 25-50 meters of relief from canyon rim to floor
- **Wall Slope**: Steep to very steep, with evidence of slumping and mass wasting
- **Steps and Knickpoints**: Multiple vertical drops or "steps" along the canyon axis, creating a staircase profile

The canyon's straight trajectory is unusual among submarine canyons, which often meander. This suggests structural control, possibly related to underlying faults or fractures in the continental shelf bedrock that guided initial canyon development.

Lower Canyon and Fan Transition

The canyon transitions into the Redondo Submarine Fan system at approximately:

- **Depth of Transition**: 400-600 meters
- **Distance Offshore**: Approximately 10-15 kilometers
- **Fan Depth Range**: 600-825 meters

- **Basin**: San Pedro Basin (northern end)



Geological Formation History

Primary Formation Period: Pleistocene Epoch

Redondo Canyon was primarily carved during the **Pleistocene Epoch** (2.6 million years ago to 11,700 years ago), a period characterized by repeated glacial-interglacial cycles. The canyon achieved most of its current morphology during glacial maxima when:

1. **Sea levels dropped dramatically** (up to 122 meters below present levels)
2. **The continental shelf was exposed**, creating a coastal plain
3. **Rivers flowed across the exposed shelf** to reach the ocean at the shelf edge
4. **Erosion and sediment transport** carved deep valleys that became submarine canyons when sea levels rose again

Geological Time Scale

- **Early Formation**: Likely initiated in the Pliocene-Pleistocene transition (2-3 million years ago)

- **Maximum Carving**: During major glacial periods, especially the Last Glacial Maximum (LGM) ~20,000 years ago
- **Modern Modification**: Continued activity during the Holocene Epoch (last 11,700 years) through turbidity currents and sediment transport

The canyon represents millions of years of geological processes, but the current morphology primarily reflects late Pleistocene to Holocene sculpting.

Pleistocene Glacial Periods and Sea Level Change

The Last Glacial Maximum (LGM)

The Last Glacial Maximum occurred approximately **20,000 years ago** and represents the most recent period when Redondo Canyon would have been actively carved by subaerial and submarine fluvial processes.

Key LGM Characteristics:

- **Global Sea Level**: 111-122 meters (364-400 feet) below present mean sea level
- **Ice Sheet Extent**: Massive Laurentide and Cordilleran ice sheets covered much of North America
- **Climate**: Significantly colder and drier than present in Southern California
- **Coastline Position**: Located far seaward of the present shoreline, potentially 10-15 kilometers west of current Redondo Beach
- **Continental Shelf**: Fully exposed as a broad, vegetated coastal plain
- **River Courses**: Extended far offshore across the exposed shelf

Sea Level Fluctuations

During the Pleistocene, sea level fluctuated dramatically through multiple glacial-interglacial cycles:

Period	Time (Years Ago)	Sea Level Relative to Present	Shelf Exposure
Last Glacial Maximum	~20,000	-111 to -122 meters	Fully exposed
Late Glacial	~15,000	-90 to -100 meters	Mostly exposed
Early Holocene	~10,000	-40 to -60 meters	Partially exposed

Period	Time (Years Ago)	Sea Level Relative to Present	Shelf Exposure
Mid-Holocene	~6,000	-5 to -10 meters	Near present
Present	0	0 meters (baseline)	Modern shelf

Exposed Continental Shelf Environment

During glacial lowstands, the exposed continental shelf off what is now Los Angeles would have been:

- A **coastal plain** extending 10-20 kilometers west of the present shoreline
- Covered with **terrestrial vegetation** adapted to a cooler, drier climate
- Traversed by **major rivers** including the ancestral San Gabriel River
- Subject to **subaerial erosion** by rainfall, runoff, and fluvial processes
- Home to **Pleistocene megafauna** including mammoths, saber-toothed cats, ground sloths, and American lions

The San Gabriel River and other drainages would have carved valleys across this plain, and these valleys continued cutting downward even as they extended offshore, creating the precursors to modern submarine canyons.

The San Gabriel River Connection

Why the San Gabriel River?

Redondo Canyon is geologically associated with the **San Gabriel River**, not the Los Angeles River, based on:

1. **Geographic Alignment**: The canyon's location and trajectory align with the paleo-drainage pattern of the San Gabriel River during lowstand conditions
2. **Sediment Volume**: The size and extent of the Redondo Fan system matches the sediment discharge capacity of a major river like the San Gabriel
3. **Regional Drainage Patterns**: During the Pleistocene, the San Gabriel River drained a vast watershed in the San Gabriel Mountains and would have delivered enormous sediment loads to this area
4. **Comparative Analysis**: Other Southern California submarine canyons (Hueneme, Mugu, Dume) show clear associations with specific river systems; Redondo's characteristics match the San Gabriel River's parameters

San Gabriel River Watershed

The San Gabriel River drains approximately **689 square kilometers** (266 square miles) of the San Gabriel Mountains and Los Angeles Basin, including:

- **High Mountain Terrain**: Peaks exceeding 2,000 meters elevation with steep slopes
- **Highly Erodible Geology**: Fractured and weathered granitic and metamorphic rocks
- **High Relief**: Over 2,000 meters of elevation change from headwaters to sea level
- **Flood-Prone**: Capable of generating massive debris flows and floods during intense storms

During the Pleistocene, the river would have been even more powerful, with:

- Greater discharge from wetter glacial-period climates in the mountains
- Higher sediment loads from periglacial erosion and mass wasting
- Extended channel length across the exposed continental shelf
- Direct connection to the developing canyon system

Pleistocene River Course

During lowstand periods, the San Gabriel River would have:

1. **Flowed westward** across the Los Angeles Basin (as it does today)
2. **Continued across the exposed continental shelf** for an additional 10-15 kilometers
3. **Carved a valley** (proto-Redondo Canyon) as it approached the shelf edge
4. **Created a shelf-edge delta** or river mouth at approximately -100 to -120 meters paleodepth
5. **Delivered sediment** directly into the developing canyon head

The river's sediment load—including sand, gravel, and finer materials—would have been funneled into the canyon, triggering turbidity currents and other mass-wasting events that further deepened and extended the canyon system.

Submarine Canyon Formation Mechanisms

Submarine canyons form through a combination of processes operating over geological time scales. Redondo Canyon's formation involved multiple mechanisms working in concert.

1. Subaerial River Erosion (Glacial Lowstands)

During glacial periods with exposed continental shelves:

- **Fluvial Incision**: The ancestral San Gabriel River cut a valley across the shelf through normal river erosion processes
- **Downcutting**: The river eroded downward into bedrock and sediment as it flowed toward the lowered sea level
- **Valley Development**: Over thousands of years, the valley deepened and widened, creating the basic canyon framework
- **Knickpoint Migration**: Waterfalls and rapids within the river system migrated upstream, creating the stepped profile visible in the modern canyon

This process is analogous to how rivers carve canyons on land (like the Grand Canyon), but operating on the continental shelf during ice age conditions.

2. Turbidity Currents (Primary Ongoing Mechanism)

Turbidity currents are the dominant process shaping submarine canyons during both lowstand and highstand conditions:

What Are Turbidity Currents?

- Dense, sediment-laden underwater flows that travel down submarine slopes like underwater avalanches
- Composed of suspended sediment (sand, silt, clay) mixed with seawater
- Driven by gravity—the dense sediment-water mixture is heavier than surrounding seawater
- Capable of traveling at velocities exceeding 70 kilometers per hour
- Can transport enormous volumes of sediment hundreds of kilometers

Turbidity Current Triggers:

- Earthquake shaking that destabilizes canyon wall sediments
- Storm waves that suspend sediment at the canyon head
- Submarine landslides and slumps on canyon walls
- River flood events that deliver massive sediment pulses
- Tidal currents and internal waves that mobilize accumulated sediment

Erosional Power: Turbidity currents erode submarine canyons through:

- **Abrasion**: Sediment particles scour the canyon floor and walls like liquid sandpaper
- **Hydraulic Plucking**: High-velocity flow tears rock fragments from the substrate
- **Dissolution**: Chemical erosion of soluble minerals
- **Undercutting**: Basal erosion undermines canyon walls, triggering slumps and failures

Modern research using seafloor sensors has documented turbidity currents in Redondo Canyon at depths of 1,300 meters, confirming that these processes remain active today.

3. Mass Wasting and Submarine Landslides

Canyon walls are inherently unstable due to:

- **Oversteepening** from turbidity current erosion
- **Sediment Accumulation** on steep slopes
- **Earthquake Shaking** in this tectonically active region
- **Fluid Overpressure** in sediments

Failure mechanisms include:

- **Slumps**: Large coherent blocks sliding downslope
- **Debris Flows**: Liquefied sediment-water mixtures
- **Rockfalls**: Individual blocks falling from canyon walls
- **Creep**: Slow, continuous downslope movement

These mass-wasting events:

- Contribute sediment to the canyon floor
- Widen the canyon by removing wall material
- Can trigger turbidity currents
- Create the irregular, stepped morphology visible in bathymetric data

4. Groundwater Seepage and Sapping

Submarine groundwater discharge can contribute to canyon formation through:

- **Seepage Erosion**: Groundwater flowing through sediment weakens canyon walls
- **Spring Sapping**: Concentrated discharge creates alcoves and undermines slopes
- **Fluid Overpressure**: Reduces sediment strength, promoting failure
- **Chemical Weathering**: Alters and weakens bedrock

While less important than turbidity currents, groundwater processes may have contributed to Redondo Canyon's development, particularly during lowstand periods when hydraulic gradients were steep.

5. Tidal and Internal Wave Processes

Modern oceanographic processes contribute to ongoing canyon modification:

- **Tidal Currents**: Oscillating flows that resuspend sediment
- **Internal Waves**: Subsurface waves that propagate along density boundaries (thermoclines)
- **Upwelling**: Canyon topography induces upwelling of cold, nutrient-rich water
- **Rip Currents**: Shore-perpendicular flows that deliver sediment to the canyon head

These processes are relatively subtle compared to turbidity currents but operate continuously, gradually modifying canyon morphology.

Turbidity Currents and Sediment Transport

Turbidity currents are the lifeblood of submarine canyons, maintaining their relief and transporting sediment from the continental shelf to the deep sea.

Anatomy of a Turbidity Current

A typical turbidity current consists of three parts:

1. **Head**: The leading edge, thickest and most concentrated, with the highest velocity
2. **Body**: The main flow, gradually decreasing in concentration and velocity downstream
3. **Tail**: The trailing dilute flow, carrying fine-grained suspended sediment

Flow Dynamics in Redondo Canyon

Initiation at Canyon Head:

- Sediment accumulates at the canyon head (10-30 meters depth) from littoral drift and storm waves
- A triggering event (earthquake, storm, critical accumulation) initiates failure
- Sediment mobilizes and mixes with seawater, forming a dense suspension

Acceleration Down-Canyon:

- The flow accelerates due to gravity as it descends the steep canyon floor
- Entrains additional sediment by eroding the canyon bed
- Velocity increases to potentially 30-70 km/h in the steepest sections
- Flow becomes self-sustaining—erosion adds sediment, increasing density and velocity

Deposition on the Fan:

- As the flow reaches the gentler slope of the submarine fan, velocity decreases
- Sediment begins to settle out, coarsest first (graded bedding)
- Flow spreads laterally across the fan surface
- Multiple flows build up thick sequences of turbidite deposits

Turbidity Current Deposits (Turbidites)

Turbidity currents leave characteristic deposits called **turbidites** that consist of:

- **Graded Bedding**: Coarse sediment at the base fining upward to mud at the top
- **Sharp Basal Contact**: Erosional surface marking the passage of the flow head
- **Sedimentary Structures**: Flame structures, load casts, ripple marks, and parallel lamination
- **Bouma Sequence**: A classical vertical succession of sedimentary structures

The Redondo Fan is composed of hundreds to thousands of individual turbidite layers deposited over the Pleistocene and Holocene, representing a long history of turbidity current activity.

Recent Measurements

In 2012, USGS researchers deployed sensors in Redondo Canyon at **1,300 meters depth** and recorded:

- **Small-scale turbidity currents** with sediment concentrations of ~0.017%
- **Flow velocities** sufficient to transport fine sand and silt
- **Associated submarine landsliding** on canyon walls
- **Continuous background transport** of suspended sediment by weaker currents

These observations confirm that Redondo Canyon remains an active sediment conduit even during the current sea-level highstand.

Littoral Drift and Modern Sediment Supply

Even though the San Gabriel River no longer flows directly into Redondo Canyon (it empties into San Pedro Bay ~20 km southeast), the canyon continues to receive sediment through **littoral drift**—the longshore transport of sediment by waves and currents.

Coastal Sediment Transport System

Southern California's coast experiences:

- **Prevailing Wave Direction**: From the northwest and west
- **Net Longshore Transport**: Predominantly southward along the coast
- **Sediment Sources**: Cliff erosion, river inputs (Santa Clara River, Ventura River, Los Angeles River system), beach sand
- **Transport Rates**: Hundreds of thousands of cubic meters of sand per year

Sediment Delivery to Redondo Canyon

Several mechanisms deliver sediment to the shallow canyon head:

1. Longshore Drift

- Sand moves southward along the beach
- Encounters the submarine canyon head just offshore
- Some sediment "falls" into the canyon, escaping the littoral system

2. Rip Currents

- Strong shore-perpendicular currents generated by wave setup
- Transport suspended sediment offshore
- Converge at the canyon head, delivering sediment directly into the canyon

3. Storm Waves

- Large winter waves (4-6 meters) suspend vast amounts of sediment
- Increased turbulence mobilizes sand from the nearshore zone
- Sediment-laden water flows into the canyon head

4. Preferential Transport Corridor (PTC)

- Recent research has identified a "preferential transport corridor" extending 10-15 km along the inner shelf toward the San Gabriel Canyon (adjacent to Redondo Canyon)
- Created by wave refraction and modified circulation due to canyon bathymetry
- Funnels sediment toward canyon heads
- Represents a significant sediment delivery pathway

Littoral Convergence

Bathymetric studies have identified **littoral convergence** above the Redondo Canyon head—a zone where:

- Longshore transport from both north and south converges
- Wave refraction over the canyon creates complex current patterns
- Enhanced sediment accumulation occurs
- Conditions favor sediment delivery to the canyon

This convergence zone explains why Redondo Canyon remains active despite the San Gabriel River's current discharge location.

Seasonal Variability

Sediment supply varies seasonally:

- **Winter (November-March)**: Large northwest swells dominate; maximum southward transport; highest sediment delivery

- **Spring (April-June)**: Transitional conditions; moderate transport
- **Summer (July-September)**: Smaller south swells; reduced or reversed transport; minimal delivery
- **Fall (October-November)**: Return to northwest swells; increasing transport

This seasonal pattern means canyon activity likely peaks in winter and early spring when sediment supply and storm wave energy are greatest.

The Redondo Submarine Fan

Submarine fans are the underwater equivalents of alluvial fans on land—depositional features where sediment accumulates at the base of a sediment conduit (river or canyon).

Redondo Fan Morphology

The Redondo Submarine Fan is a complex depositional system with:

- **Total Area**: Approximately 40-60 square kilometers
- **Depth Range**: 600-825 meters
- **Location**: Northern San Pedro Basin
- **Thickness**: Potentially hundreds of meters of turbidite sediment
- **Surface Features**: Multiple distributary channels, levees, lobes, and sediment waves

Channel-Levee Systems

The fan surface exhibits at least **5 longitudinal levee-bordered channels**, indicating:

- Multiple phases of activity with shifting channel locations
- Repeated turbidity current events that built natural levees (similar to river levees but underwater)
- Complex depositional history over thousands to millions of years

Current Activity:

- Only **one channel is currently active**
- Other channels are abandoned and partially filled
- The active channel shows fresh scours and erosional features
- Inactive channels display smooth, sediment-draped surfaces

Sediment Waves

Large **sediment waves** (dune-like features) occur on the Redondo Fan surface:

- **Wavelength**: 100-300 meters
- **Height**: 5-15 meters
- **Origin**: Created by turbidity currents through a process called "cyclic steps"
- **Significance**: Indicate high-energy turbidity currents capable of large-scale sediment transport

These features are visible on high-resolution multibeam bathymetry and indicate repeated powerful flows.

Depositional Environment

The fan accumulates sediment through:

1. **Channelized Flows**: Turbidity currents confined to active channels deposit sand in channel floors
2. **Overbank Deposition**: Flows that overtop channel levees deposit fine sediment on the fan surface
3. **Lobe Formation**: When channels terminate, flows spread and deposit sediment in lobate bodies
4. **Hemipelagic Sedimentation**: Continuous rain of fine particles from the water column between turbidity current events

The result is a thick sequence of interbedded turbidites and hemipelagic mud recording the canyon's activity over geological time.

Modern Activity and Holocene Dynamics

Despite the sea-level highstand conditions of the Holocene Epoch (last ~11,700 years), Redondo Canyon remains an active geological system.

Evidence for Continued Activity

Morphological Features:

- Fresh-looking erosional scours in channel floors
- Steep, unvegetated canyon walls (no significant marine growth indicating recent activity)
- Lack of sediment draping on active channel features
- Sharp relief maintained between canyon floor and walls

Sedimentological Evidence:

- Recent turbidite deposits in cores from the fan
- Fine-scale channel shifting within the fan's active channel system

- Multiple superimposed channel features indicating repeated flows

Direct Observations:

- Turbidity current events recorded by seafloor instruments (2012 USGS study)
- Submarine landslide scars indicating recent mass-wasting
- Measurements of suspended sediment transport

Holocene Sea-Level Rise

Sea level rose approximately **120 meters** from the Last Glacial Maximum (~20,000 years ago) to the present, drowning the continental shelf and moving the coastline landward. This had several effects:

1. **Disconnection**: The San Gabriel River no longer flows directly to the canyon head
2. **Reduced Sediment Supply**: Littoral drift provides less sediment than direct river input
3. **Decreased Activity Frequency**: Turbidity currents are less frequent than during lowstand periods
4. **Continued Maintenance**: Despite reduced activity, the canyon maintains its relief through periodic events

Many submarine canyons become inactive during highstands, but Redondo Canyon's shallow head allows it to continue intercepting shelf sediments.

Controls on Highstand Activity

Research has identified several factors that allow submarine canyons to remain active during highstands:

1. **Shallow Canyon Head**: Allows interception of littoral drift and storm-generated sediment
2. **Wave Climate**: High wave energy delivers sediment to the canyon head
3. **Structural Position**: Alignment with sediment transport pathways
4. **Tectonic Activity**: Earthquakes trigger canyon wall failures and turbidity currents
5. **Sediment Supply**: Adequate sediment delivery from coastal erosion and littoral drift

Redondo Canyon benefits from all these factors, explaining its continued activity.

California Continental Borderland Context

Redondo Canyon exists within the unique geological setting of the California Continental Borderland—a 500-kilometer-long province of submarine basins, banks, and ridges extending from Point Conception to Baja California.

Borderland Characteristics

Unlike typical passive continental margins with smooth continental shelves and slopes, the Borderland features:

- **Complex Topography**: Alternating ridges (banks) and basins
- **Tectonic Control**: Fault-bounded features related to transform plate boundary tectonics
- **Deep Basins**: Isolated basins up to 2,000 meters deep
- **Shallow Banks**: Submarine highs rising to within 100 meters of sea level
- **Limited Sediment Escape**: Basins trap sediment rather than allowing transport to the deep ocean

San Pedro Basin

Redondo Canyon delivers sediment to the **San Pedro Basin**, a structural depression with:

- **Maximum Depth**: ~900 meters
- **Area**: ~1,100 square kilometers
- **Sediment Fill**: Kilometers of accumulated turbidites and hemipelagic sediment
- **Tectonic Origin**: Bounded by the Palos Verdes Fault and other structures
- **Closed Configuration**: All sediment entering the basin is trapped

The northern end of San Pedro Basin is filled by the Redondo Submarine Fan, with contributions from other canyons as well.

Comparison with Adjacent Basins

Santa Monica Basin (to the north):

- Receives sediment from Santa Monica, Redondo, and Dume Canyons
- Completely closed basin—100% sediment retention
- Deeper than San Pedro Basin (~900+ meters)
- Anoxic bottom waters preserve detailed sediment records

Los Angeles Basin (onshore):

- Related structural province formed by similar tectonic processes
- Contains oil-bearing sedimentary rocks deposited in similar submarine environments during the Miocene-Pliocene
- Demonstrates long-term evolution of borderland tectonics

Tectonic Setting

The Borderland's complex structure results from:

1. **Transform Plate Boundary**: Pacific Plate moving northwest relative to North American Plate
2. **Transtension**: Pull-apart basins forming in zones of oblique extension
3. **Active Faulting**: Ongoing deformation creating earthquakes and topographic relief
4. **Crustal Rotation**: Some blocks rotating due to complex fault interactions
5. **Recent Origin**: Most current topography developed in the last 5-20 million years

This tectonic activity contributes to submarine canyon formation by:

- Creating steep relief favorable to mass wasting
- Triggering earthquakes that initiate turbidity currents
- Continuously modifying regional topography
- Providing accommodation space (basins) for sediment accumulation

---

Observing Redondo Canyon Today

Google Earth and Digital Bathymetry

Redondo Canyon is clearly visible on modern bathymetric visualization platforms:

Google Earth:

- Switch to terrain view and zoom to Redondo Beach, California
- The canyon appears as a dark linear feature extending southwest from the harbor
- Depth shading shows the dramatic relief
- 3D perspective view reveals the canyon's steep walls and stepped profile

NOAA Bathymetric Data Viewer:

- High-resolution multibeam sonar data available
- Shows canyon morphology in detail
- Reveals submarine fan features and channels

USGS Coastal & Marine Geology Program:

- Scientific-quality bathymetric maps and data
- GIS datasets for detailed analysis
- Published maps and reports

Modern Seafloor Mapping Technology

Recent surveys of Redondo Canyon employed:

1. **Multibeam Sonar**: Acoustic systems that map the seafloor in high resolution (1-5 meter grid spacing)
2. **Side-Scan Sonar**: Creates images of seafloor texture and features
3. **Sub-Bottom Profilers**: Penetrate sediment to reveal subsurface structure
4. **Underwater Photography**: ROVs and AUVs capture visual imagery
5. **Sediment Coring**: Samples sediment for laboratory analysis

These technologies have revolutionized our understanding of submarine canyons, revealing details invisible to earlier generations of scientists.

Recreation and Tourism

Redondo Canyon is a popular destination for:

SCUBA Diving:

- Accessible from shore at Veterans Park (Redondo Beach)
- Depths of 30-40 meters within recreational diving limits
- Known for cold water (mid-50s°F) due to upwelling
- Dramatic wall diving with steep drop-offs
- Unique marine life attracted to canyon environment

Sport Fishing:

- Canyon attracts pelagic fish species
- Upwelling brings nutrients supporting food webs
- Deep-water species accessible relatively close to shore

Educational Programs:

- Used by universities for oceanography and marine biology courses
- Public outreach highlighting submarine geology
- Citizen science opportunities

The canyon's accessibility makes it valuable for both scientific research and public education.

Comparison with Other Southern California Canyons

Southern California has numerous submarine canyons, each with unique characteristics:

Monterey Canyon (Central California)

- **Comparison Benchmark**: Often compared to the Grand Canyon in scale
- **Depth**: Over 3,600 meters deep
- **Length**: ~470 kilometers including Monterey Fan
- **River Association**: Ancestral Salinas River
- **Activity Level**: Extremely active with frequent turbidity currents
- **Unique Features**: Canyon head in Monterey Bay; deep-sea research station (MARS)

Redondo Canyon is much smaller and shallower but shares similar formation mechanisms.

Hueneme and Mugu Canyons (Ventura County)

- **River Association**: Santa Clara River (major sediment source)
- **Activity**: Highly active due to abundant sediment supply
- **Depth**: ~600 meters to basin floors
- **Characteristics**: Receive direct river discharge during floods

These canyons demonstrate the importance of direct river connections for canyon activity.

Santa Monica Canyon

- **Location**: Adjacent to Redondo Canyon, immediately to the north
- **Formation**: Associated with ancestral Los Angeles River/Ballona Creek system
- **Current Activity**: Much less active than Redondo
- **Morphology**: Smoother, more sediment-filled appearance
- **Explanation**: Less effective sediment capture compared to Redondo Canyon

The contrast between Santa Monica and Redondo Canyons highlights the importance of canyon head position and sediment supply.

La Jolla Canyon (San Diego County)

- **Depth**: Approaches 1,000 meters
- **Accessibility**: Canyon head very close to shore at La Jolla Cove
- **Geology**: Cuts through bedrock, exposing ancient rock formations
- **Recreation**: Extremely popular diving and snorkeling site

La Jolla Canyon is similar to Redondo in having a very shallow, accessible canyon head.

Newport Canyon (Orange County)

- **River Association**: Santa Ana River (Pleistocene)
- **Formation**: Carved during glacial lowstands similar to Redondo
- **Current Status**: Less active than Redondo
- **Fan System**: Deposits in San Pedro Basin

Newport Canyon represents a similar formation history but different modern activity level.

Scientific Research and Discoveries

Historical Research (1960s-1980s)

Early studies of Redondo Canyon used:

- **Fathometers**: Single-beam depth sounders creating sparse bathymetric profiles
- **Sediment Cores**: Piston and gravity cores revealing turbidite sequences
- **Geological Sampling**: Dredging and grab sampling of canyon walls
- **Submersibles**: Manned deep-sea vehicles for direct observation

Key findings included:

- Recognition of the Redondo Submarine Fan
- Identification of multiple leveed channels
- Documentation of turbidite deposits
- Association with San Gabriel River paleo-drainage

Seminal Publication: Gorsline, D.S. and Emery, K.O. (1959). "Turbidity-current deposits in San Pedro and Santa Monica Basins off Southern California" *Geological Society of America Bulletin*, provided early understanding of Southern California canyon-fan systems.

Modern Research (1990s-Present)

Recent investigations employed:

- **Multibeam Mapping**: Complete high-resolution bathymetric coverage
- **Seafloor Observatories**: Long-term instrument deployments
- **Numerical Modeling**: Computer simulations of turbidity currents
- **Repeat Surveys**: Detecting morphological changes over time

Key 21st Century Studies:

USGS Small-Scale Turbidity Current Study (2012):

- Deployed sensors at 1,300 meters depth
- Recorded small turbidity currents with 0.017% sediment concentration
- Documented associated submarine landsliding
- Demonstrated continued canyon activity

California Seafloor Mapping Program (CSMP):

- Complete multibeam coverage of California state waters
- High-resolution imagery of canyon heads and walls
- Public data availability for research and management

Submarine Canyon Dynamics Research:

- Studies of preferential transport corridors delivering sediment to canyons
- Wave-canyon interaction modeling
- Sediment wave formation by cyclic steps in turbidity currents
- Canyon influence on coastal sediment budgets

Ongoing Research Questions

Scientists continue investigating:

1. **Frequency of Large Turbidity Currents**: How often do major canyon-flushing events occur?
2. **Sediment Budget**: What percentage of littoral drift is captured by the canyon?
3. **Earthquake Triggering**: What magnitude earthquake is required to trigger turbidity currents?
4. **Climate Change Impacts**: Will changing wave climate affect sediment delivery?
5. **Stratigraphic Record**: Can fan deposits record past earthquakes and climate events?
6. **Biological Connectivity**: How does the canyon influence marine ecosystems and larval dispersal?

Ecological Significance

Beyond its geological importance, Redondo Canyon is ecologically significant:

Physical Oceanography

Upwelling:

- Canyon bathymetry modifies coastal circulation
- Induces upwelling of cold, nutrient-rich deep water
- Creates locally cool water temperatures (lower 50s°F even in summer)
- Supports enhanced primary productivity

Internal Waves:

- Generated by tidal currents interacting with canyon topography
- Mix nutrients into surface waters
- Create dynamic habitat for marine life

Biological Productivity

The canyon supports:

Planktonic Communities:

- Enhanced phytoplankton growth from upwelled nutrients
- Zooplankton concentrations attracted to productivity

Benthic (Seafloor) Communities:

- Cold-water species not typical of Southern California shallow waters
- Sponges, corals, and other filter feeders
- Specialized canyon fauna adapted to steep, hard substrates

Pelagic Fish:

- Rockfish species using canyon walls as habitat
- Juvenile fish nursery areas
- Predatory fish attracted to prey concentrations

Marine Mammals

Cetaceans (Whales and Dolphins):

- Gray whales migrate past canyon during annual migration
- Dolphins and porpoises feed in productive canyon waters
- Occasional whale sightings attributed to canyon upwelling

Pinnipeds (Seals and Sea Lions):

- California sea lions forage in canyon areas
- Harbor seals occasionally observed

Conservation Considerations

Submarine canyons are increasingly recognized as:

- **Biodiversity Hotspots**: Supporting unique assemblages of species
- **Essential Fish Habitat**: Important for commercial and recreational fisheries
- **Carbon Sequestration**: Rapidly burying organic carbon in sediments

- **Vulnerable Ecosystems**: Susceptible to human impacts (trawling, pollution, climate change)

Redondo Canyon falls within California's system of Marine Protected Areas (MPAs), providing some conservation protection.

Conclusions

Redondo Submarine Canyon is a remarkable geological feature that:

1. **Formed Primarily During Pleistocene Glacial Periods**: Carved by the ancestral San Gabriel River when sea levels were up to 122 meters lower than present, exposing the continental shelf

2. **Demonstrates Multiple Formation Mechanisms**: Created through subaerial river erosion during lowstands and submarine turbidity currents during both lowstand and highstand conditions

3. **Remains Dynamically Active**: Continues to transport sediment from the continental shelf to the deep sea through turbidity currents, mass wasting, and other processes

4. **Exhibits Complex Morphology**: Features a remarkably shallow canyon head (10-30 meters depth), steep walls, stepped profile, and terminates in an extensive submarine fan system

5. **Represents an Active Sediment Conduit**: Captures littoral drift and storm-generated sediment, funneling it through the canyon to the Redondo Fan in San Pedro Basin

6. **Is Accessible and Observable**: Visible on Google Earth and other bathymetric platforms; accessible to recreational divers; valuable for scientific research and public education

7. **Has Ecological Importance**: Creates unique oceanographic conditions supporting enhanced biological productivity and specialized marine communities

8. **Provides a Geological Archive**: Fan deposits preserve a record of canyon activity, past sea levels, climate changes, and tectonic events extending back millions of years

Broader Significance

Redondo Canyon exemplifies fundamental Earth processes:

- **Erosion and Sedimentation**: The perpetual movement of material from high elevations to low
- **Sea Level Change**: How glacial-interglacial cycles reshape continental margins
- **Tectonics**: The influence of plate boundary deformation on seafloor morphology
- **Mass Wasting**: Gravity-driven sediment transport shaping submarine landscapes

Understanding Redondo Canyon contributes to:

- **Geohazard Assessment**: Submarine landslides and turbidity currents can damage seafloor infrastructure
- **Coastal Management**: Canyon influence on beach erosion and sediment budgets
- **Climate Science**: Canyon deposits record past climate and sea level changes
- **Marine Conservation**: Protecting canyon ecosystems and their ecological functions

Future Perspectives

As research continues, Redondo Canyon will remain important for:

1. **Monitoring Submarine Processes**: Real-time observation of turbidity currents and canyon evolution
2. **Climate Change Studies**: How changing wave climate and sediment supply affect canyon activity
3. **Paleoclimate Reconstruction**: Extracting records of past conditions from fan deposits
4. **Ecological Baselines**: Understanding canyon ecosystems before further human impact
5. **Educational Outreach**: Bringing submarine geology to public awareness

Redondo Submarine Canyon stands as a testament to the dynamic nature of Earth's surface—continuously shaped by the interplay of rivers, oceans, gravity, and deep time.

Acknowledgments

This report synthesizes findings from decades of scientific research conducted by the United States Geological Survey (USGS), National Oceanic and Atmospheric Administration (NOAA), university researchers, and marine geologists. The ability to observe submarine canyons in unprecedented detail through multibeam sonar and other technologies has revolutionized our understanding of these features.

Report Compiled: February 2026
Primary Research Period: 1950s-2025

Canyon Age: ~2-3 million years (Pliocene-Pleistocene to present)
Formation Context: Pleistocene glacial lowstands and ongoing Holocene processes

---